Fluzone® High-Dose

Consumer Medicine Information (CMI) summary

The <u>full CMI</u> on the next page has more details. If you are worried about using this vaccine, speak to your doctor, nurse or pharmacist.

1. Why am I using Fluzone High-Dose?

Fluzone High-Dose is a vaccine. This vaccine is recommended to help protect against influenza (flu) and associated pneumonia in persons aged 60 years and older.

For more information, see Section 1. Why am I using Fluzone High-Dose? in the full CMI.

2. What should I know before I am given Fluzone High-Dose?

Do not use if you have ever had an allergic reaction to Fluzone High-Dose or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions or take any other medicines.

For more information, see Section 2. What should I know before I am given Fluzone High-Dose? in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Fluzone High-Dose and affect how it works.

A list of these medicines is in Section 3. What if I am taking other medicines? in the full CMI.

4. How is Fluzone High-Dose given?

Fluzone High-Dose is administered to you by a health care professional.

More instructions can be found in Section 4. How is Fluzone High-Dose given? in the full CMI.

5. What should I know about being given Fluzone High-Dose?

| Things you | Tell your doctor, nurse or pharmacist if you notice | |
|--|--|--|
| should do | signs of allergic reactions that may include difficulty breathing, shortness of breath, swelling of the face, lips, throat or tongue, cold, clammy skin, palpitations, dizziness, weakness, fainting, rash or itching. | |
| Looking after your vaccine Fluzone High-Dose is usually stored in the doctor's surgery or clinic, or at the pharmacy. How need to store Fluzone High-Dose: | | |
| | Keep it in the refrigerator, store at 2°C to 8°C. Do not freeze. | |

For more information, see Section 5. What should I know about being given Fluzone High-Dose? in the full CMI.

6. Are there any side effects?

Serious side effects can include severe allergic reactions (anaphylaxis) including swelling of the face, lips, tongue, throat or any other part of the body and which may cause difficulty in swallowing or breathing (angioedema). See your doctor immediately if you notice this.

Common local side effects include pain, redness, swelling, hardness and bruising at the injection site. Common systemic side effects include muscle aches, feeling unwell, headache, shivering, nausea and tiredness.

For more information, including what to do if you have any side effects, see Section 6. Are there any side effects? in the full CMI.

Fluzone® High-Dose

Active ingredient(s): Inactivated Trivalent Influenza Vaccine (Split Virion) Influenza Virus Haemagglutinin

Consumer Medicine Information (CMI)

This leaflet provides important information about using Fluzone High-Dose. You should also speak to your doctor, nurse or pharmacist if you would like further information or if you have any concerns or questions about using Fluzone High-Dose.

Where to find information in this leaflet:

- 1. Why am I using Fluzone High-Dose?
- 2. What should I know before I am given Fluzone High-Dose?
- 3. What if I am taking other medicines?
- 4. How is Fluzone High-Dose given?
- 5. What should I know about being given Fluzone High-Dose?
- <u>6.</u> <u>Are there any side effects?</u>
- 7. Product details

1. Why am I using Fluzone High-Dose?

Fluzone High-Dose contains the active ingredient influenza virus haemagglutinin. Fluzone High-Dose is a vaccine. Vaccines are used to help protect you against infectious diseases.

Fluzone High-Dose is recommended to help protect you against influenza (flu) and associated pneumonia in persons aged 60 years and older. Influenza (flu) is a disease caused by different strains of the influenza virus. The influenza virus can spread rapidly, especially in homes or institutions, where it can be very easy for someone to catch flu and spread the disease.

Fluzone High-Dose works by helping the body to make antibodies. These antibodies help the body to recognise the flu virus and help prevent the infection. It takes a few weeks for the body to produce these antibodies.

Fluzone High-Dose will only help to prevent flu if it is caused by one of thethree strains of virus contained in the vaccine or other strains closely related to them. Fluzone High-Dose will not prevent flu if you are incubating the disease before vaccination or if it is caused by another virus.

Previous injections of flu vaccine are unlikely to give you protection against the current strains of flu virus most common this year, so you will need to receive a vaccine against flu every year especially if you are at risk.

Your doctor will be able to recommend the best time for you to be vaccinated. Flu is present every year, even when epidemics or outbreaks are not reported.

2. What should I know before I am given Fluzone High-Dose?

Warnings

Do not use Fluzone High-Dose if:

- you are allergic to the active ingredients, or any of the ingredients listed at the end of this leaflet
- you have ever had a severe allergic reaction after getting any flu vaccine.

Always check the ingredients to make sure you can receive this vaccine.

Check with your doctor if you:

- have or have had Guillain-Barré syndrome (severe muscle weakness) after getting a flu vaccine
- have or have had an immune response problem because the immune response to the vaccine may be diminished
- have bleeding problems or bruise easily
- have an illness with a high or moderate temperature or an acute illness, the vaccination should be postponed until after you have recovered.

After vaccination, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section <u>6</u>. Are there any side effects?

3. What if I am taking other medicines?

Tell your doctor, nurse or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with Fluzone High-Dose and affect how it works.

Tell your doctor, nurse or pharmacist if you are taking medicines that may reduce your immune response such as corticosteroids (for example prednisone), medicines used to treat cancer (chemotherapy), radiotherapy or other medicines affecting the immune system.

Having other vaccines

Your doctor will advise you if Fluzone High-Dose may be given with another vaccine.

Check with your doctor, nurse or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Fluzone High-Dose.

4. How is Fluzone High-Dose given?

Fluzone High-Dose is administered by your doctor, nurse or pharmacist as a single 0.5 mL injection injected into the muscles of the upper arm.

5. What should I know about being given Fluzone High-Dose?

Things you should do

Call your doctor straight away if you notice:

 signs of allergic reactions that may include difficulty breathing, shortness of breath, swelling of the face, lips, throat or tongue, cold, clammy skin, palpitations, dizziness, weakness, fainting, rash or itching.

Driving or using machines

Do not drive or use machines if you are feeling unwell after vaccination. Wait until any effects of the vaccine have worn off before you drive or use machines.

Looking after your vaccine

Fluzone High-Dose is usually stored in the doctor's surgery or clinic, or at the pharmacy. However, if you need to store Fluzone High-Dose:

- Keep Fluzone High-Dose in the original packaging until it is time for it to be given.
- Keep it in the refrigerator, store at 2C to 8C. Do not freeze Fluzone High-Dose.

Do not use Fluzone High-Dose after the expiry date which is stated on the carton after EXP.

Do not use Fluzone High-Dose if the packaging is torn or shows signs of tampering.

Keep it where young children cannot reach it.

Getting rid of any unwanted vaccine

If you no longer need to use this vaccine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this vaccine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor, nurse or pharmacist if you have any further questions about side effects.

Less serious side effects

| Less s | erious side effects | What to do |
|--|--|------------|
| • It | ching (pruritis) | they worry |
| Syste | mic side effects: | you. |
| • m • fe • he • fe • d • cc • ve • di • in • ra • in • (c) | | you. |
| • te | emporary reduction in the umber of blood particles called latelets (thrombocytopenia) wollen glands in neck, armpit or | |
| gı | roin (lymphadenopathy) | |

Serious side effects

| Serious side effects | What to do |
|---|--|
| Severe allergic reactions (anaphylaxis) including swelling of the face, lips, tongue, throat or any other part of the body and which may cause difficulty in swallowing or breathing (angioedema) rare skin condition with severe blisters and bleeding in the lips, eyes, mouth, nose and genitals (Stevens-Johnson Syndrome) numbness or pins and needles sensation (paresthesia) neurological disorders that may result in stiff neck, confusion, numbness, pain and weakness of the limbs, loss of balance, loss of reflexes, paralysis of part or all the body (encephalomyelitis and transverse myelitis, brachial neuritis, Guillain-Barré Syndrome), fits (convulsions including febrile convulsions), facial palsy (Bell's palsy), vision | Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects. |

| Serious side effects | What to do |
|---|------------|
| disorders due to the optic nerves dysfunction (optic neuritis/neuropathy), fainting (syncope) shortly after vaccination • blood vessel inflammation (vasculitis) which may result in skin rashes and in very rare cases in temporary kidney problems, blood vessel opening (vasodilatation) • chest pain • wheezing, throat tightness, difficulty breathing (dyspnoea) • excess of blood in the white of the eye (ocular hyperaemia). | |

Tell your doctor, nurse or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this vaccine.

Always make sure you speak to your doctor, nurse or pharmacist before you decide to stop taking any of your medicines.

7. Product details

What Fluzone High-Dose contains

| Active ingredient (main ingredient) | Influenza virus haemagglutinin of the following strains A/Victoria/4897/2022 (H1N1)pdm09-like strain (A/Victoria/4897/2022, IVR-238) A/Thailand/8/2022 (H3N2)-like strain (A/California/122/2022, SAN-022) B/Austria/1359417/2021-like strain (B/Michigan/01/2021, wild type) |
|--|--|
| Other ingredients (inactive ingredients) | sodium chloride, dibasic sodium phosphate, monobasic sodium phosphate, octoxinol-9, water for injection, and traces of formaldehyde. |

| Potential allergens | Fluzone High-Dose contains less |
|---------------------|---------------------------------|
| | than 1 microgram ovalbumin |
| | (egg protein) per dose. |

Do not receive this vaccine if you are allergic to any of these ingredients.

Fluzone High-Dose does not contain any antibiotics or preservative.

Fluzone High-Dose syringes are not made with natural rubber latex.

What Fluzone High-Dose looks like

Fluzone High-Dose is supplied as single-dose, 0.5mL prefilled syringe without needles (AUST R 473685)

Who distributes Fluzone High-Dose

Distributed in Australia by: sanofi-aventis australia pty ltd 12 - 24 Talavera Road Macquarie Park NSW 2113

Australia

Freecall: 1800 818 806

Email: medinfo.australia@sanofi.com

This leaflet was prepared in November 2025.

fluz-hd-ccdsv4-cmiv1-21nov25