

GLICLAZIDE MR VIATRIS

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about taking this medicine, speak to your doctor or pharmacist.

1. Why am I taking GLICLAZIDE MR VIATRIS?

GLICLAZIDE MR VIATRIS contains the active ingredient gliclazide. GLICLAZIDE MR VIATRIS is used to control blood glucose (sugar) in patients with Type II diabetes mellitus. For more information, see Section [1. Why am I taking GLICLAZIDE MR VIATRIS?](#) in the full CMI.

2. What should I know before I take GLICLAZIDE MR VIATRIS?

Do not take if you have ever had an allergic reaction to gliclazide or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, are pregnant or breastfeeding or plan to become pregnant or breastfeed. For more information, see Section [2. What should I know before I take GLICLAZIDE MR VIATRIS?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with GLICLAZIDE MR VIATRIS and affect how it works. A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I take GLICLAZIDE MR VIATRIS?

Your doctor will tell you how many tablets to take; swallow them whole with water at the same time each day, with breakfast.

More instructions can be found in Section [4. How do I take GLICLAZIDE MR VIATRIS?](#) in the full CMI.

5. What should I know while taking GLICLAZIDE MR VIATRIS?

Things you must do	<ul style="list-style-type: none">• Tell your doctors, dentists, pharmacists and diabetes educators that you are taking this medicine.• Make sure that you, your friends, family and work colleagues can recognise the symptoms of hypoglycaemia and hyperglycaemia and know how to treat them.• If you become pregnant while taking GLICLAZIDE MR VIATRIS, tell your doctor immediately.• Make sure you check your blood glucose levels regularly to tell if your diabetes is being controlled.
Things you must not do	<ul style="list-style-type: none">• Do not skip meals while taking GLICLAZIDE MR VIATRIS.• Do not give GLICLAZIDE MR VIATRIS to anyone else, even if they have the same condition as you.• Do not stop or change your GLICLAZIDE MR VIATRIS dose without checking with your doctor.
Driving or using machines	<ul style="list-style-type: none">• GLICLAZIDE MR VIATRIS may cause dizziness and drowsiness in some people.• GLICLAZIDE MR VIATRIS may lower your blood glucose levels may slow your reaction time and affect your ability to drive or operate machinery.
Drinking alcohol	<ul style="list-style-type: none">• If you drink alcohol while taking GLICLAZIDE MR VIATRIS, you may get flushing, sensation of warmth, headache, breathing difficulties, rapid heartbeat, stomach pains or feel sick and vomit.
Looking after your medicine	<ul style="list-style-type: none">• Keep them in a cool dry place below 25°C. Protect from moisture.• Keep your tablets in the pack until it is time to take them.

For more information, see Section [5. What should I know while taking GLICLAZIDE MR VIATRIS?](#) in the full CMI.

6. Are there any side effects?

Serious side effects: skin rash, redness itching and/or hives, angioedema (rapid swelling of tissues such as eyelids, face, lips, mouth, tongue or throat that may result in breathing difficulty). Common side effects: low or high blood sugar, cold-like or respiratory symptoms (runny/blocked nose, sneezing, sinus or throat discomfort, coughing, bronchitis, upper respiratory infection), back or joint pain, high blood pressure, chest pain, headache, weakness, viral or urinary infections, dizziness, stomach upset (nausea, pain, vomiting, diarrhoea, constipation), or skin blistering/peeling. For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

GLICLAZIDE MR VIATRIS

Active ingredient(s): *gliclazide*

Consumer Medicine Information (CMI)

This leaflet provides important information about taking GLICLAZIDE MR VIATRIS. **Speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about taking GLICLAZIDE MR VIATRIS.**

Where to find information in this leaflet:

1. [Why am I taking GLICLAZIDE MR VIATRIS?](#)
2. [What should I know before I take GLICLAZIDE MR VIATRIS?](#)
3. [What if I am taking other medicines?](#)
4. [How do I take GLICLAZIDE MR VIATRIS?](#)
5. [What should I know while taking GLICLAZIDE MR VIATRIS?](#)
6. [Are there any side effects?](#)
7. [Product details](#)

1. Why am I taking GLICLAZIDE MR VIATRIS?

GLICLAZIDE MR VIATRIS contains the active ingredient gliclazide. Gliclazide belongs to a group of medicines called sulphonylureas. The medicine releases the active ingredient gliclazide progressively over 24 hours.

It lowers blood glucose by increasing the amount of insulin (a hormone that controls blood glucose levels) produced by your pancreas.

GLICLAZIDE MR VIATRIS is used to control blood glucose (sugar) in patients with Type II diabetes mellitus. This type of diabetes is also known as non-insulin-dependent diabetes (NIDDM), or maturity-onset diabetes.

GLICLAZIDE MR VIATRIS is used for Type II diabetes mellitus when diet and exercise are not enough to control your blood glucose.

GLICLAZIDE MR VIATRIS can be used alone or together with insulin or other medicines for treating diabetes.

How it works

Glucose is used by the body as fuel, and all people have glucose circulating in their blood. In diabetes, levels of blood glucose are higher than is needed, which is also known as hyperglycaemia. If your blood glucose is not properly controlled, you may experience hypoglycaemia (low blood glucose) or hyperglycaemia (high blood glucose). High blood glucose can lead to serious problems with our heart, circulation and/or kidneys.

It is very important to control high blood glucose whether or not you feel unwell. This really helps to avoid serious long-term health problems, which can involve the heart, eyes, circulation, and/or kidneys.

As with many medicines used for the treatment of diabetes, there is a possibility that blood glucose levels

may become very low during treatment with GLICLAZIDE MR VIATRIS. This is known as hypoglycaemia.

Section [5. What should I know while taking GLICLAZIDE MR VIATRIS?](#) of this leaflet contains advice about recognising and treating hypoglycaemia.

Ask your doctor if you have any questions about why GLICLAZIDE MR VIATRIS has been prescribed for you.

GLICLAZIDE MR VIATRIS is not addictive.

2. What should I know before I take GLICLAZIDE MR VIATRIS?

Warnings

Do not take GLICLAZIDE MR VIATRIS if:

- **you are allergic to:**
 - o gliclazide, or any of the ingredients listed at the end of this leaflet
 - o other sulphonylureas,
 - o GLICLAZIDE MR VIATRIS related medicines such as sulfa antibiotics or to thiazide diuretics (a type of "fluid" or "water" tablet)Symptoms of an allergic reaction to GLICLAZIDE MR VIATRIS or to these medicines may include:
 - o skin rash, itchiness or hives,
 - o shortness of breath, swelling of the face, lips or tongue,
 - o muscle pain or tenderness or joint pain.

If you are not sure if you have an allergy to GLICLAZIDE MR VIATRIS, check with your doctor.

Always check the ingredients to make sure you can take GLICLAZIDE MR VIATRIS.

- **you have Type 1 diabetes mellitus (also known as insulin dependent diabetes (IDDM), or juvenile-onset diabetes)**
- **you have unstable diabetes**
- **you have diabetic ketoacidosis (a problem which affects the acidity of your blood and can lead to coma - which is mainly associated with Type 1 diabetes)**
- **you have severe kidney disease**
- **you have severe liver disease**
- **you are taking an antibiotic medicine containing the active ingredient miconazole**
- the expiry date printed on the pack has passed

If you take it after the expiry date has passed, it may not work as well.
- the packaging is torn or shows signs of tampering.

If it is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking GLICLAZIDE MR VIATRIS, talk to your doctor.

Check with your doctor if you:

- are allergic to any of the ingredients listed at the end of this leaflet; to any other medicines; or to any other substances, such as foods, preservatives or dyes
- **have a family history of or know you have the hereditary condition glucose-6- phosphate dehydrogenase (G6PD) deficiency (abnormality of red blood cells), lowering of the haemoglobin level and breakdown of red blood cells (haemolytic anaemia) can occur**
- **have had any of the following medical conditions:**
 - o kidney disease
 - o liver disease
 - o a history of diabetic coma
 - o adrenal, pituitary or thyroid problems
 - o heart failure
- take any medicines for any other condition
- **have any medical condition, or do anything, that may increase the risk of hyperglycaemia - for example:**
 - o you are ill or feeling unwell (especially with fever or infection)
 - o you are injured
 - o you are having surgery
 - o you are taking less GLICLAZIDE MR VIATRIS than prescribed
 - o you are taking on less exercise than normal
 - o you are eating more carbohydrate than normal.
 - o drinking alcoholic drinks
 - o not eating regular meals
 - o taking on more exercise than usual.

If you have not told your doctor, pharmacist or diabetes educator about any of the above, tell them before you start taking GLICLAZIDE MR VIATRIS.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Do not take GLICLAZIDE MR VIATRIS if are pregnant or trying to become pregnant.

GLICLAZIDE MR VIATRIS may affect your developing baby if you take it during pregnancy. Insulin is more suitable for controlling blood glucose during pregnancy. Your doctor will usually replace GLICLAZIDE MR VIATRIS with insulin while you are pregnant.

Do not take GLICLAZIDE MR VIATRIS are breastfeeding or plan to breastfeed.

GLICLAZIDE MR VIATRIS is not recommended while you are breastfeeding. It is not known whether GLICLAZIDE MR VIATRIS passes into breast milk.

Use in children

Do not give GLICLAZIDE MR VIATRIS to a child.

There is no experience with the use of GLICLAZIDE MR VIATRIS in children.

Use in older people

Elderly people can generally use GLICLAZIDE MR VIATRIS safely.

There are no special instructions for older people taking GLICLAZIDE MR VIATRIS.

3. What if I am taking other medicines?

Tell your doctor, pharmacist or diabetes educator if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Taking GLICLAZIDE MR VIATRIS may change the effect of some medicines, and some medicines may affect how well GLICLAZIDE MR VIATRIS works.

Some medicines may lead to low blood glucose (hypoglycemia) by increasing the blood-glucose-lowering effect of GLICLAZIDE MR VIATRIS. These include:

- some medicines used to treat fungal or yeast infections (miconazole which is contraindicated)
- alcohol
- other medicines used to treat diabetes (such as biguanides and insulin)
- some medicines used to treat high blood pressure and other heart conditions (including angiotensin receptor blocker, beta-blockers)
- some medicines used to treat depression and other mental illness (MAOIs)
- some cholesterol-lowering medicines (clofibrate)
- some medicines used to treat arthritis, pain and inflammation (including high dose aspirin, ibuprofen, phenylbutazone)
- some antibiotics (chloramphenicol; clarithromycin; sulphonamides)
- some medicines used to treat acid reflux and stomach ulcers
- some medicines used to treat gout (probenecid)
- some medicines used to treat tuberculosis.

Some medicines may lead to high blood glucose levels (hyperglycaemia) by weakening the blood glucose-lowering effect of GLICLAZIDE MR VIATRIS. These include:

- alcohol
- some medicines for epilepsy (danazol)
- some medicines used to treat depression and other mental illness (chlorpromazine)
- St John's Wort (*Hypericum perforatum*) preparations used to treat depression
- some hormones used in hormone replacement therapy and oral contraceptives (oestrogen, progesterone)
- some medicines for asthma (salbutamol, intravenous terbutaline)
- barbiturates, medicines used for sedation
- glucocorticoids
- some medicines for fluid retention (thiazides, furosemide)
- nicotinic acid.

Some medicines may lead to unstable blood glucose (low blood sugar and high blood sugar) when taken at the same time as GLICLAZIDE MR VIATRIS, especially in elderly patients. These include:

- a class of antibiotics called fluoroquinolones.

GLICLAZIDE MR VIATRIS may change the effects of some other medicines. These include:

- some medicines used to prevent blood clots (warfarin).

You may need different amounts of your medicine or you may need to take different medicines. Your doctor, pharmacist or diabetes educator can tell you what to do if you are taking any of these medicines. They also have a more complete list of medicines to be careful with or avoid while taking GLICLAZIDE MR VIATRIS.

Ask your doctor or pharmacist if you are not sure if you are taking any of these medicines.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect GLICLAZIDE MR VIATRIS.

4. How do I take GLICLAZIDE MR VIATRIS?

How much to take

- Follow all directions given to you by your doctor, pharmacist, and diabetes educator carefully.
They may differ from the information contained in this leaflet.
- If you do not understand the instructions on the box, ask your doctor or pharmacist for help.
- Your doctor will tell you how many tablets to take each day. They may increase or decrease the dose, depending on your blood glucose levels.

When to take GLICLAZIDE MR VIATRIS

- Swallow the tablets whole with a glass of water. GLICLAZIDE MR VIATRIS tablets should not be crushed or chewed.
Crushing or chewing the tablets may change the effectiveness of the tablet.
- GLICLAZIDE MR VIATRIS tablets do not have a break bar and should not be broken in half.
- It is important to take your GLICLAZIDE MR VIATRIS at the same time each day - usually with breakfast. Taking GLICLAZIDE MR VIATRIS with food can help to minimise the risk of hypoglycaemia.
- Do not skip meals while taking GLICLAZIDE MR VIATRIS.

How long to take GLICLAZIDE MR VIATRIS

- Continue taking GLICLAZIDE MR VIATRIS for as long as your doctor recommends. Make sure you keep enough GLICLAZIDE MR VIATRIS to last over weekends and holidays.

- GLICLAZIDE MR VIATRIS will help control your diabetes but will not cure it. Therefore, you may have to take it for a long time.

If you forget to take GLICLAZIDE MR VIATRIS

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as you remember (with food), then go back to taking your tablets as you would normally.

Missed doses can cause high blood glucose (hyperglycaemia).

If you are not sure whether to skip the dose, talk to your doctor or pharmacist.

Do not take a double dose to make up for the dose you missed.

If you double a dose, this may cause low blood glucose (hypoglycaemia).

If you take too much GLICLAZIDE MR VIATRIS

If you think that you or anyone else has taken too much GLICLAZIDE MR VIATRIS, urgent medical attention may be needed.

You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**) for advice, or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

Do this even if there are no signs of discomfort or poisoning.

If you take too much GLICLAZIDE MR VIATRIS together with other medicines for diabetes or alcohol, you may experience symptoms of low blood glucose (hypoglycaemia).

If not treated quickly, these symptoms may progress to loss of co-ordination, slurred speech, confusion, loss of consciousness and fitting.

At the first signs of hypoglycaemia, raise your blood glucose quickly by following the instructions at the end of this leaflet. Section [5. What should I know while taking GLICLAZIDE MR VIATRIS?](#) contains advice about recognising and treating hypoglycaemia.

If you experience any of these symptoms, immediately get medical help.

5. What should I know while taking GLICLAZIDE MR VIATRIS?

Things you must do

- **If you become pregnant while you are taking GLICLAZIDE MR VIATRIS, tell your doctor.**
- Tell all doctors, dentists, pharmacists and diabetes educators who are involved with your treatment that you are taking GLICLAZIDE MR VIATRIS.

- If you are about to start taking any new medicines, tell your doctor and pharmacist that you are taking GLICLAZIDE MR VIATRIS.
- **Take GLICLAZIDE MR VIATRIS exactly as your doctor has prescribed. Otherwise you may not get the full benefits from treatment.**
- **Make sure you check your blood glucose levels regularly.**

This is the best way to tell if your diabetes is being controlled properly. Your doctor or diabetes educator will show you how and when to do this.

- **Make sure that you, your friends, family and work colleagues can recognise the symptoms of hypoglycaemia and hyperglycaemia and know how to treat them.**

Instructions at the end of this leaflet. Section [5. What should I know while taking GLICLAZIDE MR VIATRIS?](#) can help you with this.

- **Visit your doctor regularly so that they can check on your progress.**
- **Carefully follow your doctor's and dietician's advice on diet, drinking alcohol and exercise.**
- **Tell your doctor immediately if you notice the return of any symptoms of hyperglycaemia that you had before starting GLICLAZIDE MR VIATRIS.**

These may include lethargy or tiredness, headache, thirst, passing large amounts of urine and blurred vision. These may be signs that GLICLAZIDE MR VIATRIS is no longer working, even though you may have been taking it successfully for some time.

Remind any doctor, dentist or pharmacist you visit that you are taking GLICLAZIDE MR VIATRIS.

Things you must not do

- **Do not give GLICLAZIDE MR VIATRIS to anyone else, even if they have the same condition as you.**
- **Do not take GLICLAZIDE MR VIATRIS to treat other complaints unless your doctor tells you to.**
- **Do not stop taking GLICLAZIDE MR VIATRIS, or change the dosage, without checking with your doctor.**
- **Do not skip meals while taking GLICLAZIDE MR VIATRIS.**

Things to be careful of

- **Protect your skin when you are in the sun, especially between 10am and 3pm.**

Sulphonylureas (the group of medicines that GLICLAZIDE MR VIATRIS belongs to) may cause your skin to be more sensitive to sunlight than it is normally. Exposure to sunlight may cause a skin rash, itching, redness, or a severe sunburn.

- **If outdoors, wear protective clothing and use a SPF 30+ sunscreen. If your skin does appear to be burning, tell your doctor immediately.**
- **If you are travelling, it is a good idea to:**

- o wear some form of identification showing you have diabetes
- o carry some form of sugar to treat hypoglycaemia if it occurs, for example, sugar sachets or jelly beans
- o carry emergency food rations in case of a delay, for example, dried fruit, biscuits or muesli bars
- o keep GLICLAZIDE MR VIATRIS readily available

- **If you become sick with a cold, fever or flu, it is very important to continue taking GLICLAZIDE MR VIATRIS, even if you fell unable to eat your normal meal. If you have trouble eating solid food, use sugar-sweetened drinks as a carbohydrate substitute or eat small amounts of bland food.**

Your diabetes educator or dietician can give you a list of foods to use for sick days.

Recognising and treating hyPOglycaemia (very LOW blood sugar levels)

Hypoglycaemia may occur during GLICLAZIDE MR VIATRIS treatment.

The first signs of hypoglycaemia are usually weakness, trembling or shaking, sweating, lightheadedness, dizziness, headache or lack of concentration, irritability, tearfulness, hunger, and/ or numbness around the lips and tongue.

At the first signs of hypoglycaemia take some sugar to raise your blood sugar level quickly.

Do this by eating 5 to 7 jelly beans, 3 teaspoons of sugar or honey, drinking half a can of non-diet soft drink, taking 2-3 glucose tablets or a tube of glucose gel.

Then take some extra carbohydrates -

such as plain biscuits, fruit or milk - unless you are within 10-15 minutes of your next meal. Taking this extra carbohydrate will help to prevent a second drop in your blood glucose level.

If not treated quickly, hypoglycaemia symptoms may progress to loss of co-ordination, slurred speech, confusion, fits or loss of consciousness.

If hypoglycaemia symptoms do not get better straight away after taking sugar then go to the Accident and Emergency department at your nearest hospital - if necessary by calling an ambulance.

Contact your doctor or diabetes educator for advice if you are concerned about hypoglycaemia.

Recognising and treating hyPERglycaemia (HIGH blood sugar levels)

Some people may feel fine when their glucose levels are high. Others notice symptoms of hyperglycaemia like tiredness, lack of energy, thirst, passing large amounts of urine, headache, and/or blurred vision.

If you notice symptoms of hyperglycaemia, or your blood sugar levels are high, tell your doctor immediately. You may need adjustments of the dose or type of medicines you are taking.

It is very important to control high blood glucose whether or not you feel unwell. This really helps to avoid serious

long-term health problems, which can involve the heart, eyes, circulation, and/or kidneys.

If you experience any of the signs of hyperglycaemia (high blood glucose) contact your doctor or diabetes educator for advice immediately.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how GLICLAZIDE MR VIATRIS affects you.

Also, be especially careful not to let your blood glucose levels fall too low.

GLICLAZIDE MR VIATRIS may cause dizziness and drowsiness in some people. Low blood glucose levels may also slow your reaction time and affect your ability to drive or operate machinery. Section [5. What should I know while taking GLICLAZIDE MR VIATRIS?](#) contains advice about recognising and treating hypoglycaemia.

Drinking alcohol can make this worse. If either of these occurs, do not drive, operate machinery or do anything else that could be dangerous.

Drinking alcohol

Tell your doctor if you drink alcohol.

If you drink alcohol while taking GLICLAZIDE MR VIATRIS, you may get flushing, sensation of warmth, headache, breathing difficulties, rapid heartbeat, stomach pains or feel sick and vomit.

Looking after your medicine

- Keep them in a cool dry place below 25°C. Protect from moisture.
- Keep your tablets in the pack until it is time to take them.

GLICLAZIDE MR VIATRIS will not keep as well outside its blister packaging.

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Heat and dampness can destroy some medicines.

Keep it where young children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Getting rid of any unwanted medicine

If you no longer need to take this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not take this medicine after the expiry date.

6. Are there any side effects?

If you do not feel well while you are taking GLICLAZIDE MR VIATRIS then tell your doctor, pharmacist, or diabetes educator as soon as possible.

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

However if you do - or if you notice anything else that is making you feel unwell - you should consult your doctor or pharmacist.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none">• hypoglycaemia and hyperglycaemia. Section 5. What should I know while taking GLICLAZIDE MR VIATRIS? contains advice about recognising and treating hypoglycaemia and hyperglycaemia• runny or blocked nose, sneezing, facial pressure or pain, bronchitis, sore throat and discomfort when swallowing, upper respiratory infection, coughing• back pain, arthralgia, arthrosis• high blood pressure, chest pain• headache, unusual weakness• viral infection, urinary tract infection• dizziness• stomach upset with symptoms like feeling sick, stomach pain, vomiting, diarrhoea or constipation• decrease in the number of cells in the blood (e.g. platelets, red and white blood cells) which may cause paleness, prolonged bleeding, bruising, sore throat and fever have been reported. These symptoms usually vanish when the treatment is discontinued• increase of some hepatic enzymes levels, and exceptionally a liver disease• your vision may be affected for a short time especially at the start of treatment. This effect is due to changes in blood sugar levels.	Speak to your doctor if you have any of these less serious side effects.

Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none">skin rash, redness itching and/or hives, angioedema (rapid swelling of tissues such as eyelids, face, lips, mouth, tongue or throat that may result in breathing difficulty) have been reported. The rash may progress to widespread blistering or peeling of the skin and may be the first sign of rare life-threatening conditions (e.g. Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and severe hypersensitivity reactions (DRESS). Exceptionally, DRESS have been reported: initially as flu-like symptoms and a rash on the face then an extended rash with a high temperatureas for other sulphonylureas, the following adverse events have been observed: cases of severe changes in the number of blood cells and allergic inflammation of the wall of blood vessels, reduction in blood sodium (hyponatraemia), symptoms of acute pancreatitis, symptoms of liver impairment (e.g. jaundice) which in most cases disappeared after withdrawal of the sulphonylurea, but may lead to life-threatening liver failure in isolated cases.	Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.

Tell your doctor, pharmacist or diabetes educator if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What GLICLAZIDE MR VIATRIS contains

Active ingredient	gliclazide 30 mg
-------------------	------------------

(main ingredient)	
Other ingredients (inactive ingredients)	hypromellose magnesium stearate microcrystalline cellulose

Do not take this medicine if you are allergic to any of these ingredients.

What GLICLAZIDE MR VIATRIS looks like

GLICLAZIDE MR VIATRIS 30 mg tablets are white to off-white, oblong, biconvex tablet debossed with 'M' on one side of the tablet and 'GL 30' on the other side (AUST R 295541).

They are available in blister packs of 100 tablets.

Who distributes GLICLAZIDE MR VIATRIS

Alphapharm Pty Ltd trading as Viatris

Level 1, 30 The Bond

30-34 Hickson Road

Millers Point NSW 2000

www.viatris.com.au

Phone: 1800 274 276

This leaflet was prepared in December 2025.

GLICLAZIDE MR VIATRIS_cmi\Dec25/00