Consumer Medicine Information (CMI) summary

The <u>full CMI</u> on the next page has more details. If you are worried about using IMFINZI, speak to your doctor or pharmacist.

1. Why am I using IMFINZI?

IMFINZI contains the active ingredient durvalumab. IMFINZI is used to treat several kinds of cancers.

For more information, see Section 1. Why am I using IMFINZI? in the full CMI.

2. What should I know before I use IMFINZI?

Do not use if you have ever had an allergic reaction to IMFINZI or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section 2. What should I know before I use IMFINZI? in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with IMFINZI and affect how it works.

A list of these medicines is in Section 3. What if I am taking other medicines? in the full CMI.

4. How do I use IMFINZI?

 IMFINZI will be given to you as a liquid infusion into your vein and your doctor will decide how many treatments you need.

More instructions can be found in Section <u>4. How do I use IMFINZI?</u> in the full CMI.

5. What should I know while using IMFINZI?

Things you should do	 Remind any doctor, dentist or pharmacist you visit that you are using IMFINZI.
	 Keep all of your doctor's appointments so that your progress can be checked.
Things you should not do	Do not miss a dose of IMFINZI

Driving or using machines

 Be careful before you drive or use any machines or tools until you know how IMFINZI affects you.

For more information, see Section <u>5. What should I know while using IMFINZI?</u> in the full CMI.

6. Are there any side effects?

Like all medicines, IMFINZI can cause side effects, although not everybody gets them. Your doctor will discuss these with you and will explain the risks and benefits of your treatment.

When you take IMFINZI, you can have some serious side effects.

For more information, including what to do if you have any side effects, see Section <u>6. Are there any side</u> effects? in the full CMI.

Active ingredient(s): durvalumab

Consumer Medicine Information (CMI)

This leaflet provides important information about using IMFINZI. You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using IMFINZI.

Where to find information in this leaflet:

- 1. Why am I using IMFINZI?
- 2. What should I know before I use IMFINZI?
- 3. What if I am taking other medicines?
- 4. How do I use IMFINZI?
- 5. What should I know while using IMFINZI?
- 6. Are there any side effects?
- 7. Product details

1. Why am I using IMFINZI?

IMFINZI contains the active ingredient durvalumab.

IMFINZI is a monoclonal antibody, a type of protein. It is a type of immunotherapy and belongs to a group of medicines called immune checkpoint inhibitors. It works with your immune system to destroy cancer cells.

IMFINZI can be used to treat a type of lung cancer called non-small cell lung cancer (NSCLC). It may be prescribed to you:

- in combination with chemotherapy that contains platinum prior to surgery (neoadjuvant treatment) and alone after surgery (adjuvant treatment) if:
 - your cancer has spread within your lung and is able to be removed by surgery

• alone if:

- your cancer has spread within your lung and cannot be removed by surgery, and
- you have tried radiotherapy and chemotherapy that contains platinum, and your cancer has shrunk or has not worsened.

IMFINZI can be used to treat a type of lung cancer called small cell lung cancer (SCLC). It may be prescribed to you:

• alone if:

- you have limited stage SCLC (LS-SCLC) and
- your cancer has not been removed by surgery and
- your cancer has responded or stabilised after initial treatment with radiotherapy and chemotherapy that contains platinum.

• in combination with chemotherapy if:

- you have extensive stage SCLC (ES-SCLC) where your cancer has spread within your lungs (or to other parts of the body) and
- you have not received previous treatment.

IMFINZI in combination with chemotherapy can be used to treat a type of cancer called biliary tract cancer (BTC), such as cancer of the bile ducts (cholangiocarcinoma) and gallbladder. It may be prescribed to you if:

 your cancer has spread within these regions (or to other parts of the body).

IMFINZI in combination with tremelimumab can be used to treat a type of liver cancer, called unresectable hepatocellular carcinoma (uHCC). It may be prescribed to you if your cancer:

- cannot be removed by surgery (unresectable), and
- has spread within your liver (or to other parts of the body)

IMFINZI can be used to treat a type of cancer of the lining of the uterus (endometrial cancer) that has spread beyond the original tumour or come back (recurred). It may be prescribed to you in combination with chemotherapy, followed by IMFINZI alone.

A test may be used to find out the mismatch repair (MMR) status of your endometrial cancer to help your doctor decide which treatment option may be the best for your cancer.

IMFINZI is used to treat a type of bladder cancer called muscle-invasive bladder cancer (MIBC). It is used in combination with chemotherapy prior to the surgical removal of your bladder (neoadjuvant treatment) followed by IMFINZI alone after surgery (adjuvant treatment). It may be prescribed to you if:

 your cancer has spread into the muscle layer of the bladder but not to other parts of the body

When IMFINZI is given in combination, it is important that you also read the Consumer Medicine Information (CMI) for the specific anti-cancer medicines you may be receiving. If you have any questions about these medicines, ask your doctor.

Your doctor may have prescribed IMFINZI for another reason.

2. What should I know before I use IMFINZI?

Warnings

Do not use IMFINZI if:

- you are allergic to durvalumab, or any of the ingredients listed at the end of this leaflet.
- Always check the ingredients to make sure you can use IMFINZI.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

Check with your doctor if you:

- have an autoimmune disease (a condition where the body attacks its own cells) like Crohn's disease, ulcerative colitis or lupus. If you already have an autoimmune disease, your risk of immune-mediated side effects may be higher with immune checkpoint inhibitor medicines (such as IMFINZI) as they may cause inflammation in parts of your body. You may also experience more frequent flares of your autoimmune disease, which in the majority of cases are mild
- have had an organ transplant
- have lung or breathing problems
- have liver problem
- have any other medical conditions
- take any medicines for any other condition.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section <u>6</u>. Are there any side effects?

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

You should not be given IMFINZI if you are pregnant.

It may affect your developing baby if you use it during pregnancy.

If you could become pregnant, you must use adequate birth control while you are being treated with IMFINZI and for at least 3 months after your last dose.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

You should not be given IMFINZI if you are breastfeeding.

It is not known if the active ingredient in IMFINZI passes into breast milk, but if it does, there is a possibility that your baby may be affected.

Do not breast-feed if you are being given IMFINZI and for at least 3 months after the last dose.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect IMFINZI.

4. How do I use IMFINZI?

How much and when to take IMFINZI

 IMFINZI will be given to you as a liquid infusion into your vein (IV). An infusion takes about 1 hour and will

normally be given every 2, 3 or 4 weeks. Depending on your type of cancer, IMFINZI may be given in combination with other anti-cancer medicines.

Your doctor will decide how many treatments you need.

When IMFINZI is given on the same day as chemotherapy, the IMFINZI infusion goes first, followed by chemotherapy.

When IMFINZI is given on the same day as tremelimumab, the tremelimumab infusion goes first, followed by IMFINZI.

Please refer to the CMI of the other anti-cancer medicines in order to understand the use of these other medicines. If you have questions about these medicines, ask your doctor.

If you miss an appointment to be given IMFINZI

Call your doctor right away to reschedule your appointment. It is very important that you do not miss a dose of IMFINZI.

5. What should I know while using IMFINZI?

Things you should do

Keep all of your doctor's appointments so that your progress can be checked.

Remind any doctor, dentist or pharmacist you visit that you are using IMFINZI.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how IMFINZI affects you.

IMFINZI is not expected to affect your ability to drive a car or operate machinery.

Looking after your medicine

- Store unopened vials under refrigeration at 2°C to 8°C in the original carton to protect from light.
- Do not freeze.
- Do not shake.

Follow the instructions in the carton on how to take care of your medicine properly.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use IMFINZI or it is out of date, take it to any pharmacy for safe disposal.

Do not use IMFINZI after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

IMFINZI can cause your immune system to attack normal organs and tissues in many areas of your body and can affect the way they work and this can cause the side effects.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

When IMFINZI is given in combination with other anticancer medicines (eg chemotherapy or tremelimumab), it is important that you also read the Consumer Medicine Information (CMI) for the specific anti-cancer medicines you may be receiving. If you have any questions about these medicines including side-effects, ask your doctor.

Serious side effects

```
Seriou What
      to do
side
effects
(signs
and
symptoms)
Proble Call
with
      your
      doctor
your
lungs: straight
• lung away
      or go
  infla
      straight
  that
      to
  may
      the
  inclu
       Emergency
  new
       Department
  or
      at
  word
      your
  coug
       nearest
  shor
       hospital
  of
       if
  brea
      you
  ches
      notice
  pain
      any
chro
      of
  obst
       these
  pulm serious
  diso
```

```
Seriou What
side
      to do
effects
(signs
and
symptoms)
  (CO side
  whid effects.
  caus Your
  the
      doctor
  narre
      may
  of
 airw give
 and you
      other
  mak
      medicines
  it
      in
  diffid
      order
  to
      to
  brea
      prevent
• lung more
  infed severe
  (pne complications
  or
      and
  influ reduce
  that your
  may symptoms.
  incly Your
  coud doctor
  of
      may
  phle withhold
  feve the
  chill next
```

Seriou What side to do effects (signs and symptoms) and dose diffid of brea IMFINZI Proble or stop with your your treatment liver: with • infla IMFINZI. of the liver that may inclu naus or vom feeli less hung pain on the right side

Seriou What to do side effects (signs and symptoms) of stom yelld of skin or white of eyes drow dark urine or blee or bruis more easi than norn Proble with your intesti

SeriousWhat side to do effects (signs and symptoms)

inflat of the intes may inclu diarr or more bow mov than usua blac tarry stick stoo or stoo with bloo or muc

seve

stom

pain

Serious What to do side effects (signs and symptoms) or tend Proble with some of your hormo glands • infla of your horn glan may inclu head that will not go awa or unus head

Seriouside effects	to do
_	
(signs	
and	
sympt	oms)
0)/1/4	
extre	
tired	
weig	
gain	
or	
weig	
loss,	
dizzi	
or	
faint	
feeli	
more	
hung	
or	
thirs	
than	
usua	
hair	
loss,	
feeli	
cold	
cons	
char	
to	
your	
voice	

Serious What to do side effects (signs and symptoms) urina more ofter than usua naus or vom ston area (abd pain char in moo or beha such as decr sex drive incre anxi irrita or

Seriou What to do side effects (signs and symptoms) forge S, fast and deep brea conf or a swe sme to your brea a swee or meta taste in your mou or a diffe odou

IMFINZI®

to

Serious What to do side effects (signs and symptoms) your urine or swe Proble with your kidney • inflai of your kidn that may inclu char in the amo or colo of your urine swel

Seriou What to do side effects (signs and symptoms) in your ankl or loss of appe Proble with your urinar tract: infed that may inclu the need to urina urge and frequ burn pain

Serious What to do side effects (signs and symptoms) or sens whe urina blad still feels full after urina pain abov your pubi bone bloo in the urine Proble with your skin or mouth

Seriou What to do side effects (signs and symptoms) • infla of the skin or mou may inclu rash itchii skin bliste or ulce in the mou or othe muc men • thrus in the mou

Seriouside effects (signs and sympt	to do	
Proble with your nose and throat		
nose or throa infector(sinuor tons)		
Proble with your heart: • inflat		n
of your hear may inclu ches pain		

Seriouside effects (signs and sympt	to do	
shor of brea or irreg hear Proble with your muscl		
• inflatof your mus may inclu mus weal tired and/ or pain and/ or rapid		n

Seriou What to do side effects (signs and symptoms) fatig of the mus in one or more area of your body **Proble** with your blood low num of bloo plate may inclu blee

Seriou What to do side effects (signs and symptoms) (eg nose or gum blee and/ or bruis Infusio related reaction may inclu chills or shak itchii or rash flush shor of brea or whe

Serious What to do side effects (signs and symptoms) dizzi feve feeli like pass out, back or neck pain or facia swel Encep or menin may inclu seizi neck stiffr head feve chills vom

	<u> </u>
Seriouside effects (signs and sympt	to do
eye sens to light conf and slee Guilla Barré syndre	
may incluing pain weal and para in the extre	
Gener body:	
an inflation inflationrespondingthat	

Serious What to do side effects (signs and symptoms) may inclu low or high body temp incre hear rate, incre brea rate and/ or abno white bloo cell cour Proble with your immui syster

SeriousWhat side to do effects (signs and symptoms)

• imm syste attac your red bloo cells sign and sym may inclu unus weal and fatig with incre hear rate and brea diffid

IMFINZI®

yelld

of

Seriou What to do side effects (signs and symptoms) the skin or white of the eyes (jaur dark urine and/ or an enla sple Inflam of the spinal cord: sym may inclu pain num

Serious What to do side effects (signs and symptoms) tingli or weal in the arm or legs blad or bow prob inclu need to urina more frequ urina inco e, diffid urina and cons

```
Seriou What
      to do
side
effects
(signs
and
symptoms)
Proble Contact
with
      a
     medical
your
eyes: eye
• infla (ophthalmologist)
 your straight
  of
 eyes away,
      or go
  (uve
      straight
      to
  sign
      the
  and
      Emergency
  sym
      Department
  inclu
      at
  eye
 redn your
      nearest
  eye
      hospital
  pain
      if
  light
 sens you
      notice
  and/
      these
  or
      side
  char
      effects
  in
      as
  visio
      you
```

Seriou	ı \$ What	
side	to do	
effects	5	
(signs		
and		
sympt	oms)	
	may	
	need	
	urgen	t
	medic	al
	attenti	on.

Immune checkpoint inhibitor class effects

There have been cases of the following serious side effects reported during treatment with other medicines in this class which may also occur during treatment with IMFINZI:

- decreased ability of the pancreas to make digestive enzymes, which may result in symptoms such as diarrhoea with loose and oily stools, weight loss, metabolic bone disease, and vitamin or mineral deficiencies (known as exocrine pancreatic insufficiency)
- decreased production of blood cells, which may result in symptoms such as weakness, shortness of breath, dizziness, fast or irregular heartbeat, frequent infections, unexplained or easy bruising, or bleeding from gums or other tissues (known as aplastic anaemia)

- a condition where the immune system makes too many infection-fighting cells which may result in symptoms such as an enlarged liver and/or spleen, skin rash, lymph node enlargement, breathing problems, easy bruising, kidney abnormalities and heart problems (known as haemophagocytic lymphohistiocytosis)
- coeliac disease which may result in symptoms such as stomach pain, diarrhoea and bloating after consuming gluten-containing foods)

Call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.

Most common side effects reported in clinical trials:

Most comm side effects	
IMFINATIONE:	•
• coug	your doctor if
stom painskin	you have any
rash	of these

C S	lost omm ide ffects		
•	or itching feve swell of the legs upper traction in feet under thyroglan that can	they worry you.	
•	tired or weig gain (hyp over thyro glan whice		dism)

may

What Most commato to side effects caus man diffe sym inclu weig loss hand trem rapid or irreg hear IMFIN Speak in to combi your doctor with platin if based you chem have y [lung any cance of • naus these side • cons effects • feeli and tired they

C	lost omm ide ffects	What on to	
	or wea	worry you.	
	(fatio	you	
•	skin		
	rash		
	and		
	itchii		
	decr in		
	appe		
•	hair		
	loss		
•	diarr		
•	infla		n
	of		
	the		
	nerv		
	num		
	wea		
	tingl		
	or burn		
	pain		
	of		
	the		
	arms		

Most comm side effects	What who to	
and legs couge vom under thyroglan that can caus tired or weigh gain (hyposis)		dism)
with chemo (etopo and carbo or cispla [lung	doctor if you have any of these side	y S

What Most commato to side effects • hair and loss they • decr worry appe you. • feeli tired or weal (fatio cons vom coug IMFIN Speak to with cheme your (gemc doctor and if cispla you [biliar have tract any cance of and/ these side or bladd effects cance and they

C Si	lost omm ide ffects	What colors	
•	low num of red bloo cells (and low num of white bloo cells (neu lymp and/ or leuke low	worry you.	a, a
•	of bloo plate (thro abno kidn		topenia)

What Most commato to side effects func tests diarr stom pain cons naus vom • feeli tired or wea (fatio feve abno liver func tests decr in appe • skin rash and itchii

What Most commodo to side effects • unde thyr glan that can caus tired or weig gain (hyp dism) infla of the nerv caus num weal tingl or burn pain of the arm and legs

What Most commato to side effects bloo cells (neu and leuk • cons low num of plate openia) (thrq diarr vom stom pain decr appe • swe of legs • unde thyro glan that can caus

What Most commodo to side effects tired or weig gain (hyp dism) • abno liver func tests • mus pain • infla of the nerv caus num weal tingl or burn pain of the arm and

legs

What Most commato to side effects coug prod coug hair loss Skin rash or itchii IMFIN Speak in to combi your doctor with treme if b [hepat you ar cance have any unde of thyr these glan side that effects can and caus they tired worry or weig you. gain

What Most commato to side effects stom pain diarr abno pand func test swel of the legs feve abno liver func tests coug skin rash or itchil

Some side effects can only be found when your doctor does blood tests from time to time to check your progress (eg abnormal thyroid gland function test (underactive or overactive), high calcium level, low sodium level, low number of white blood

cells, low number of red blood cells, low number of blood platelets, abnormal liver function tests (eg aspartate aminotransferase (AST) increased; alanine aminotransferase (ALT) increased) or pancreas function tests or kidney function tests (eg blood creatinine increased)).

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell. Other less common side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of IMFINZI.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

IMFINZI is only available with a doctor's prescription.

What IMFINZI contains

Active ingredient (main ingredient)	durvalumab
-------------------------------------	------------

Other ingredients (inactive ingredients)	histidine histidine hydrochloride monohydrate
	trehalose dihydrate polysorbate 80 water for injections

Do not take IMFINZI if you are allergic to any of these ingredients.

What IMFINZI looks like

IMFINZI concentrated solution for infusion is a clear to opalescent, colourless to slightly yellow liquid in a glass vial.

Australian Registration Number(s):

IMFINZI, 500 mg (500 mg/10mL) in 10 mL vial for intravenous infusion - AUST R 283216

IMFINZI, 120 mg (120 mg/2.4mL) in 10 mL vial for intravenous infusion - AUST R 283215

Who distributes **IMFINZI**

AstraZeneca Pty Ltd ABN 54 009 682 311 66 Talavera Road MACQUARIE PARK NSW 2113

Telephone: 1800 805 342

This leaflet was prepared 1 October 2025

IMFINZI is a trade mark of the AstraZeneca group of companies.

© AstraZeneca 2025

VV-RIM-01430609 v24