

MenQuadfi®

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about receiving this vaccine, speak to your doctor, nurse or pharmacist.

- ▼ This medicine is new or being used differently. Please report side effects. See the [full CMI](#) for further details

1. Why am I or my child receiving MenQuadfi?

MenQuadfi is a vaccine. It is used to help protect you or your child against infections caused by a bacteria (germs) called “Neisseria meningitides” types A, C, W and Y.

For more information, see Section [1. Why am I or my child receiving MenQuadfi?](#) in the full CMI.

2. What should I know before I or my child receive MenQuadfi?

Do not use if you or your child have ever had an allergic reaction to MenQuadfi or any of the ingredients listed at the end of the CMI. **Talk to your doctor, nurse**

or pharmacist if you or your child have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before my child or I receive MenQuadfi?](#) in the full CMI.

3. What if I or my child is taking other medicines?

Some medicines may interfere with MenQuadfi and may affect how it works. Tell your doctor nurse or pharmacist if you or your child are taking, have recently taken or might take any other vaccines or medicines, including medicines obtained without a prescription.

For more information, see Section [3. What if I or my child is taking other medicines?](#) in the full CMI.

4. How is MenQuadfi given?

MenQuadfi is given by your doctor, nurse or pharmacist.

More instructions can be found in Section [4. How is MenQuadfi given?](#) in the full CMI.

5. What should I know while being given MenQuadfi?

<p>Things you should do</p>	<p>Call your doctor, nurse or pharmacist straight away if you:</p> <ul style="list-style-type: none">• notice signs of allergic reaction which include rash, itching, difficulty breathing, shortness of breath or swelling of the face, lips, throat or tongue. <p>Driving or using machines</p> <ul style="list-style-type: none">• MenQuadfi is not likely to affect your ability to drive or use machines. However, do not drive or use any machines if you are feeling unwell.
<p>Looking after your vaccine</p>	<p>MenQuadfi is usually stored in the doctor's surgery or clinic, or at a pharmacy. However, if you need to store MenQuadfi: keep in the fridge between 2-8°C. Do not freeze.</p>

For more information, see Section [5. What should I know while being given MenQuadfi?](#) in the full CMI.

6. Are there any side effects?

Serious side effects can include allergic reactions. See your doctor immediately if you notice this. In infants 6 weeks - 12 months of age, common side effects include tenderness, redness, swelling or bruising at the injection site, feeling irritable, crying, loss of appetite, feeling drowsy, fever and vomiting. In children 12-23 months of age, common side effects include tenderness, redness or swelling at the injection site, feeling irritable, crying, loss of appetite, feeling drowsy, fever, vomiting and diarrhoea. Common side effects in children (2 years of age and older), adolescents, adults and elderly include pain, redness, or swelling at the injection site, muscle pain, headache, generally feeling unwell and fever.

For more information, including what to do if you or your child have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information.

You can help by reporting any side effects you may get. You can report side effects to your doctor, or directly at www.tga.gov.au/reporting-problems.

MenQuadfi®

Active ingredient(s): *Meningococcal Polysaccharides Groups A, C, Y and W-135*

Consumer Medicine Information (CMI)

This leaflet provides important information about receiving MenQuadfi. **You should also speak to your doctor, nurse or pharmacist if you would like further information or if you have any concerns or questions about being given MenQuadfi.**

This vaccine can be given to adults and children so you may be reading this leaflet for you or for your child.

Where to find information in this leaflet:

- [1. Why am I or my child receiving MenQuadfi](#)
 - [2. What should I know before I or my child receive MenQuadfi?](#)
 - [3. What if I or my child is taking other medicines?](#)
 - [4. How is MenQuadfi given?](#)
 - [5. What should I know while being given MenQuadfi?](#)
-

[6. Are there any side effects?](#)

[7. Product details](#)

1. Why am I or my child receiving MenQuadfi

MenQuadfi contains the active ingredients meningococcal polysaccharide groups A, C, W and Y.

MenQuadfi is a vaccine. It is used to help protect you or your child against infections caused by a bacteria (germs) called “Neisseria meningitides” types A, C, W and Y.

Neisseria meningitidis is passed from person to person and can cause meningitis, an inflammation of the tissues that surround the brain and spinal cord, or septicaemia, an infection of the blood. Both can result in serious disease with lasting effects and possibly death.

MenQuadfi can be given to infants (from 6 weeks of age), children, adolescents, adults and the elderly.

MenQuadfi stimulates the body’s natural defenses (immune system), to produce its own protection (antibodies) against the bacteria (“Neisseria meningitidis” types A, C, W and Y) that may cause the meningococcal disease. However, as with all vaccines, 100% protection cannot be guaranteed.

MenQuadfi will not prevent meningitis (an infection of the brain and spinal cord coverings) caused by other groups of Neisseria meningitidis.

2. What should I know before I or my child receive MenQuadfi?

Warnings

MenQuadfi must not be given:

- if you or your child are allergic to the active ingredients or any of the ingredients listed at the end of this leaflet. Symptoms of allergic reaction may include rash, itching, difficulty breathing, shortness of breath or swelling of the face, lips, throat, or tongue. If you are not sure if you or your child are allergic, talk to your doctor, nurse or pharmacist before you or your child receive MenQuadfi.

Always check the ingredients to make sure you or your child can receive this vaccine.

Tell your doctor, nurse or pharmacist if you or your child have:

- an infection with high temperature (over 38°C). If this applies, the vaccination will be given after the infection is under control. There is no need to delay vaccination for a minor infection such as a cold. However, talk to your doctor, pharmacist or nurse first
- a disease weakening your body's natural defences (immune system). The response to vaccines may not be optimal.
- ever fainted from an injection, Fainting sometimes with falling, can occur (mostly in adolescents and

young children) during, following, or even before, any injection with a needle.

After vaccination, you or your child may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

If you are pregnant, think you may be pregnant, plan to become pregnant or are breast-feeding, you must tell your doctor, nurse or pharmacist, before receiving MenQuadfi.

3. What if I or my child is taking other medicines?

Some medicines may interfere with MenQuadfi and may affect how it works. Tell your doctor, nurse or pharmacist if

- You or your child are taking, have recently taken or might take any other vaccines or medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.
- You or your child have been treated with medicines that may affect the body's natural defenses (immune system) such as high-dose corticosteroids or chemotherapy. The response to vaccines may not be optimal.

For age appropriate situations, MenQuadfi can be given at a separate injection site during the same visit with other vaccines. Your doctor will advise you if MenQuadfi is to be given at the same time as another vaccine.

Check with your doctor, nurse or pharmacist if you are not sure about what medicines, vitamins or supplements you or your child are taking and if these affect MenQuadfi.

4. How is MenQuadfi given?

How much is given

MenQuadfi is given by your doctor, nurse or pharmacist as a 0.5 mL injection in the muscle in the upper arm or in the thigh depending on your age and muscle mass.

5. What should I know while being given MenQuadfi?

Call your doctor straight away if you:

Notice signs of allergic reaction which include rash, itching, difficulty breathing, shortness of breath or swelling of the face, lips, throat or tongue.

Driving or using machines

MenQuadfi is not likely to affect your ability to drive or use machines. However, do not drive or use any machines if you are feeling unwell.

Looking after your vaccine

MenQuadfi is usually stored in the doctor's surgery or clinic, or at the pharmacy. However, if you need to store MenQuadfi:

- keep it where young children cannot reach it.
- keep MenQuadfi in the original pack until it is time for it to be given.
- keep it in the refrigerator, store at 2°C to 8°C. Do not freeze MenQuadfi.

Do not use MenQuadfi after the expiry date which is stated on the carton after EXP.

Do not use MenQuadfi if the packaging is torn or shows signs of tampering.

Getting rid of any unwanted medicine

Medicines including vaccines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Are there any side effects?

All medicines including vaccines can have side effects. If you or your child do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor, nurse or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<p>In infants 6 weeks to 12 months of age</p> <ul style="list-style-type: none">• Tenderness, redness, swelling or bruising where the injection was given• Feeling irritable• Crying• Loss of appetite• Feeling drowsy• Fever• Vomiting <p>In children 12 to 23 months of age</p>	<p>Speak to your doctor if you or your child have any of these less serious side effects and they worry you.</p>

Less serious side effects	What to do
<ul style="list-style-type: none"> ● tenderness, redness or swelling where the injection was given ● feeling irritable ● crying ● loss of appetite ● feeling drowsy ● fever ● vomiting ● diarrhoea <p>In persons 2 years of age and older</p> <ul style="list-style-type: none"> ● pain, redness, swelling, itching, warmth, bruising or rash where the injection was given ● muscle pain ● headache ● feeling unwell ● fever ● vomiting ● feeling dizzy ● nausea ● fatigue 	

Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none">● Signs of allergic reaction:<ul style="list-style-type: none">○ rash○ itching○ difficulty breathing,○ shortness of breath○ swelling of the face, lips, throat, or tongue● Fits (convulsions) with or without fever	<p>Call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Tell your doctor, nurse or pharmacist if you notice anything else that may be making you or your child feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you or your child experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems or in New Zealand at <https://pophealth.my.site.com/carmreportnz/s/>. By reporting side effects, you can help provide more information on the safety of this vaccine.

7. Product details

This vaccine is only available with a doctor's prescription.

What MenQuadfi contains

Active ingredients (main ingredients)	10 micrograms of each meningococcal polysaccharide groups A, C, Y and W-135 joined to approximately 55 micrograms of tetanus toxoid
Other ingredients (inactive ingredients)	Sodium chloride, sodium acetate and water for injections

Do not receive this vaccine if you or your child are allergic to any of these ingredients.

What MenQuadfi looks like

MenQuadfi is a clear, colourless solution for injection.

MenQuadfi is available in packs of 1 or 10 single dose (0.5 mL) vials. Not all pack sizes may be marketed.

AUST R 325682

Who distributes MenQuadfi

Distributed in Australia by:

sanofi-aventis australia pty ltd

12-24 Talavera Road

Macquarie Park NSW 2113

Freecall: 1800 818 806

Email: medinfo.australia@sanofi.com

Distributed in New Zealand by:

Pharmacy Retailing (NZ) Ltd t/a Healthcare Logistics

PO Box 62027

Sylvia Park Auckland 1644

Freecall: 0800 283 684

Email: medinfo.australia@sanofi.com

This leaflet was prepared in March 2026.

menq-ccds13-cmiv6-30mar26